

Infekcije maternice u mlijekočnih krava - update



Goran Bačić – Klinika za porodništvo i reprodukciju
e-mail: bacic@gef.hr

Infekcije maternice

Akutni metritis

Kronični endometritis

Subklinički endometritis

Pijometra

- Infekcija maternice je jedna od najčešćih poremetnji u puerperiju
- Nagradno pitanje: Koje su ostale dvije?

Akutni metritis

Karakteristike:

- Unutar 10 dana od poroda
- Crvenkasto smeđi vodenasti smrđljivi iscijedak
- Temp. \geq od 39.5°C
- Promjena općeg stanja - depresija
- Smanjen apetit i uzimanje hrane
- Smanjena proizvodnja mlijeka

Akutni metritis

Bakterijske kulture:

- *Escherichia coli*
- *Arcanobacterium pyogenes*
- Obligatni anaerobi - *Fusobacterium necroforum* i *Prevotella* spp.

Akutni metritis

Terapija:

- Sistemski atb bez intrauterine manipulacije
- Najbolji su *Ceporex™*, *Cephalexin®* i *Excenel®*

Kronični endometritis

Karakteristike:

- Pojava purulentnog ili mukopurulentnog iscjetka nakon više od 3 tjedna postpartum
- Temperatura obično nije povišena
- Za dijagnozu bolje vaginalno nego rektalno
- Lošija reprodukcija
- Blagi pad mlijecnosti

Kronični endometritis

Terapija:

- PGF 2α
- Cephapirin (Excenel®) intrauterino
- Proteolitički enzimi
- Sve je to dubioza jer rezultati istraživanja nisu konzistentni
- Pitanja???

AKUTNI METRITIS

Postpartum metritis

Toksični puerperalni metritis

Septični metritis



Fig. 10-4 Uterus; cow. Postpartum endometritis.

AKUTNI METRITIS

Rizični faktori za nastanak AM

- 1) maternica
- 2) metaboličke poremetnje
- 3) ravnoteža između patogenosti i imunosti

AKUTNI METRITIS

1) Maternica

- Distocije
- Mrtvorodjeni
- Blizanci
- Carski rez
- Zaostala posteljica
- Producena involucija maternice

AKUTNI METRITIS

2) Metaboličke poremetnje

- Mlijecna groznica - hipokalcemija
- Ketoza
- Dislokacija sirišta

AKUTNI METRITIS

3) Ravnoteža između patogenosti i imunosti

- Funkcija neutrofila - zakazala
- Tip bakterijske flore
- Terapija glukokortikoidima i progesteronom
- Rano stvaranje žutog tijela
- Nivo higijene

AKUTNI METRITIS

INFEKCIJA MATERNICE U
POSTPARTALNOM PERIODU I
POSLJEDIČNA UPALA MORA SE
PRIHVATITI KAO FIZIOLOŠKI
PROCES

- Preko 90% svih krava u puerperiju ima infekciju maternice
- Oko 70% prođe bez terapije

AKUTNI METRITIS

- Najčešći izolati bakterija *E. coli*,
Arcanobact. pyogenes i obligatni
anaerobi (*Fusobact. Necroforum* i
Prevotella spp.)
- Jačinu infekcije uvjetuje:
 - Količina i patogenost bakterija
 - Efikasnost mehanizma maternične obrane

AKUTNI METRITIS

Mehanizam maternične obrane

1) Anatomske i fizičke barijere

- Vulva
- Cerviks

2) Stanični i humoralni imuni sustav

- Inicijalni stanični odgovor na infekciju - pojava PMNL i makrofaga u materničnoj stijenci
- Kasnije se otpuštaju imunoglobulini i opsonini iz endometrija

AKUTNI METRITIS


Terapija:

- Posljednjih godina - sistemski ATB
- Penicilin, oksitetraciklini i cefalosporini
- Mnoštvo istraživanja - različiti rezultati i uspjeh liječenja.
- Ceftiofur dostiže MIC₉₀ za *A. pyogenes*, *E. coli* i *F. necroforum* u krvi i endometriju samo 2 sata nakon aplikacije.

AKUTNI METRITIS

Terapija:


- Cephalexin® doza: 1ml/25kg (7mg/kg/dan) tijekom 5 dana
- karenca 4 dana za meso
- za mlijeko nema



AKUTNI METRITIS

Terapija:

- Ceporex Injection™ doza: 1ml/25kg (7mgkg/dan) tijekom 5 dana
- karenca 4 - 15 dana za meso
- za mlijeko nema



Akutni metritis

Terapija:

- Excenel® RTU
- 2ml/50kg 5dana
- Karenca - nema

<http://www.freshcow.com/>



There is a smarter approach to fresh cow management

Pfizer, we've dedicated ourselves to helping dairy producers maximize the effectiveness of their fresh cow management, by helping to pioneer many of the protocols that now define a good fresh cow program, and by introducing innovative products like EXCENEL® RTU (ceftiofur hydrochloride Sterile Suspension).

EXCENEL RTU allows you to keep cows in the milking string and minimize the negative effects of cow movement and hospital pen commingling, thanks to its unique advantage of zero milk discard and proven efficacy for treating metritis. To learn how EXCENEL RTU can help you achieve a smarter approach to fresh cow management, talk to your veterinarian or Pfizer representative today. For the latest fresh cow information and resources, please visit www.freshcow.com.

With all drugs, EXCENEL RTU should not be used in animals
and to be hypersensitive to the product.



Pfizer Animal Health



EXCENEL® RTU
(ceftiofur hydrochloride
Sterile Suspension)

AKUTNI METRITIS

Terapija:

- Oksitetraciklini i penicilini
- Izbjegavati zbog slabijeg efekta i duge karence

AKUTNI METRITIS

Terapija:

- Za mlijekočna goveda jedini oksitetraciklin koji je odobren za upotrebu je:

Pfizer Animal Health US Dairy Product Overview

LIQUAMYCIN® LA-200® (oxytetracycline) Injectable

Description

LIQUAMYCIN® LA-200® (oxytetracycline injection) is a sterile, ready-to-use solution for the administration of the broad-spectrum antibiotic oxytetracycline ([TERRAMYCIN®](#)) by injection. [TERRAMYCIN](#), discovered by Pfizer scientists, is an antimicrobial agent that is effective in the treatment of a wide range of diseases caused by susceptible gram-positive and gram-negative bacteria.

Approved Uses



AKUTNI METRITIS

Terapija:

- Lokalni ATB - oblete i otopine
- Kontroverzni rezultati istraživanja
- Nema razlike kod tretmana ZP samo sa sistemskim ATB ili sistemska + lokalni
- Negativna interakcija ATB - maternična flora
- Inhibicija materničnih obrambenih mehanizama
- Upitna djelotvornost kod upale endometrija i jajovoda

AKUTNI METRITIS

Terapija:

- Aplikacija većih doza ATB od propisanih kao i aplikacija suboptimalnih doza
- Veće - opasnost od rezidua u hrani
- Manje - razvoj rezistentnih sojeva
- Idealno - prema uputama proizvođača

AKUTNI METRITIS

Dodatna terapija:

- Nesteroidni protuupalni lijekovi – nema razlike sa i bez njih
- Estradiol – stimulacija motiliteta i imunosti maternice – nedozvoljeno u EU, u USA može
- Nema pozitivnih efekata
- Kod jakih upala i jako poremećenog općeg stanja (toksemije) infuzije ili sondiranje mogu pomoći “isprati toksine”

Pitanja?

KRONIČNI ENDOMETRITIS

- Naziva se i klinički endometritis
- Prisustvo purulentnog ili mukopurulentnog eksudata u vagini 3 i više tjedana nakon poroda
- Nema povisene temperature i lošeg općeg stanja

KRONIČNI ENDOMETRITIS

Dijagnostika:

- Adspekcija vulve, perineuma i repa
- Pregled vagine spekulom
- Rektalna pretraga
- Bris maternice
- Biopsija maternice

KRONIČNI ENDOMETRITIS

Dijagnostika - klinički nalaz rektalno:

- Asimetrični maternični rogovi
- Zadebljala maternična stijenka
- Moguće palpirati fluktuaciju tekućine u rogu
- Zadebljani cerviks

KRONIČNI ENDOMETRITIS

Dijagnostika - klinički nalaz vaginalno:

- Cerviks promjera većeg od 7.5 cm
- Zacrvenjen naročito ventralni dio
- Iscjedak koji se cijedi iz cerviksa
- Zacrvenjena vagina
- Prisutnost iscjetka u vagini

KRONIČNI ENDOMETRITIS

Dijagnostika:

- Rektalna i vaginalna pretraga su indirektne metode dijagnostike i njima se ne može procijeniti stupanj upale same maternice
- Bris maternice - nije za rutinsku primjenu jer je dosta skup u kompliciran za izvođenje
- Pozitivan nalaz bakterija potvrđuje postojanje infekcije, ali ne i endometritis

KRONIČNI ENDOMETRITIS

Dijagnostika:

- Biopsija endometrijuma - pokazuje lokalne histološke promjene na endometiju
- Nije preporučljivo jer može utjecati na plodnost

KRONIČNI ENDOMETRITIS

Dijagnostika - ocjena (scoring)

- Blagi endometritis: zamućeni iscjetak, bistra sluz s krpicama gnoja lagano povećana ili nepromijenjena maternica
- Jaki endometritis: povećana maternica ispunjena tekućim sadržajem, povećan cerviks i purulentan iscjetak
- Pijometra: jako povećana maternica ispunjena tekućinom bez vidljivog iscjetka s izraženim CL

[http://www.rvc.ac.uk/AboutUS/
Staff/sheldon/ResearchInterest
s/Gallery/index.cfm](http://www.rvc.ac.uk/AboutUS/Staff/sheldon/ResearchInterest_s/Gallery/index.cfm)

KRONIČNI ENDOMETRITIS

Dijagnostika – optimalno vrijeme

- Zbog velikog broja slučajeva samoizlječenja optimalno vrijeme dijagnostike je 3-4 tjedna nakon poroda

KRONIČNI ENDOMETRITIS

Utjecaj na reproduksijsku sposobnost

- Smanjuje se koncepcija
- Povećan broj osjemenjivanja
- Produženo vrijeme do prvog osjemenjivanja
- Produženo vrijeme od telenja do koncepcije

Povećan broj izlučenja

KRONIČNI ENDOMETRITIS

Terapija - 2 osnovne strategije

1) Intrauterina primjena ATB

2) Tretman sa PGF 2α

Pitanje: Da li uopće tretirati blage oblike?

KRONIČNI ENDOMETRITIS

Klinička ispitivanja subkliničkih endometritisa govore u prilog tome da bi se svi slučajevi trebali tretirati

KRONIČNI ENDOMETRITIS

Terapija s PGF 2α i analogima

- Bazira se na luteolitičkom djelovanju PGF 2α
- Poticanju novog ciklusa (estrusa)
- Kontrakcijama miometrija koje potiskuju patološki sadržaj maternice kroz poluotvoreni cerviks
- Porast koncentracije PMN leukocita u sluznici maternice – pojačano stvaranje imunoglobulina

KRONIČNI ENDOMETRITIS

Terapija s intrauterinim ATB

- Jednak ili bolji uspjeh od PGF 2α
- Tretirati samo one koji ne mogu s PGF 2α
- Izbjeći rezidue u mesu i mlijeku
- Izbjeći stvaranje rezistentnih sojeva

KRONIČNI ENDOMETRITIS

Terapija intrauterinim antisepticima

- Izbjegavati
- Nema pozitivnih efekata na reprodukciju
- Uništava epitel jajovoda
- Smanjuje efikasnost obrambenih organizama u maternici

Biljni ekstrakti i proteolitički enzimi - obećavaju ali još nedovoljno potvrđeni

Pitanja?

SUBKLINIČKI ENDOMETRITIS

- Odsutnost kliničkih znakova kroničnog endometritisa i povećanja materničnog lumena i stijenke maternice, ali ipak postoji endometritis

SUBKLINIČKI ENDOMETRITIS

Dijagnostika

- Ultrazvuk - mala količina tekućine u lumenu maternice
- Lažno pozitivni nalaz - estrus - provjeriti jajnike

SUBKLINIČKI ENDOMETRITIS

Dijagnostika

- Endometrialna citologija -ispranje uterusa ili uzimanje uzorka s cytobrush
- Postotak PMN u preparatu dokazuje subklinički endometritis
- Granična vrijednost (threshold) PMN je od 5-18% (različite studije razni autori)

SUBKLINIČKI ENDOMETRITIS

- Važno!!!
 - Krave sa subkliničkim endometritisom imaju smanjenu reproduktivnu sposobnost
 - Preporučuju se tretmani s pgf 2α ili intrauterini ATB (cephapirin)
 - U nekim studijama nema razlike

Izazov za veterinara

- Prava dijagnoza upale maternice
- Efikasna terapija
- Praćenje efikasnosti terapije kroz postotak ozdravljenja i reproduktivnu sposobnost

- Krajnji cilj je prevencija upala maternice pravilnom prehranom, higijenom, uvjetima kod telenja i pažljivim stručnim pomaganjem kod poroda

PIJOMETRA

- Nakupljanje gnojnog sadržaja u maternici
- Klinički nalaz vaginalno - može biti mukopurulentnog sadržaja u vagini
- Klinički nalaz rektalno - povećana maternica i CL na jednom jajniku
- Lažno pozitivan nalaz - graviditet, neoplazije
- Terapija: PGF 2α - estrus

Pijometra

Pitanja????



Najčešće infekcije maternice u ostalih domaćih životinja i kućnih ljubimaca

Pijometra



Fig. 10-11 Uterus; mare. Pyometra.

CEM - Contagious Equine Metritis

- Zarazni metritis kobila
- Akutna, jako kontagiozna spolna zaraza konja (eksperimentalno magarci)
- Profuzni mukopurulentni vaginalni iscjedak
- Inficirani pastusi i kronične infekcije kobila
- bez vanjskih znakova
- Javlja se u Evropi
- Uzročnik je *Taylorella equigenitalis*, gram - bakterija

CEM - Contagious Equine Metritis

- Prenosi se isključivo spolnim aktom
- Moguće i zaraženim instrumentima i priborom
- Izvor zaraze su najčešće inficirani pastusi – svaki skok – zaražena kobila
- Klinički znaci: mukopurulentni isjedak iz vagine 10 – 14 dana nakon pripusta, obično skraćen estrusni ciklus

CEM - Contagious Equine Metritis




Fig. 10-8 Vestibule; mare. Contagious equine metritis.
(Courtesy AC Asbury, Gainesville, Florida.)



Fig. 10-9 External genitalia; mare. Contagious equine metritis.
(Courtesy AC Asbury, Gainesville, Florida.)

CEM - Contagious Equine Metritis

- Isjedak nestaje nakon par dana, ali zaraza ostaje
- Obično ne ostaju gravidne
- Ako ostane mogu inficirati ždrijebe za vrijeme ili ubrzo nakon poroda
- Takvo ždrijebe nosi uzročnika sve do seksualne zrelosti

CEM - Contagious Equine Metritis

- Razvija se edem i hiperemija endometrija, cerviksa i sluznice vagine
- Dijagnostika – izolacija uzročnika
- Bris endometrija – po mogućnosti u estrusu
- Također bris iz klitoralne fose i sinusa
- Kod pastuha – uretralna fosa, uretra, prepucij, preejakulatorna tekućina ili ejakulat

CEM - Contagious Equine Metritis

- Pastusi moraju pokazati najmanje 3 negativna rezultata da bi se proglašili CEM free i dobili licencu za rasplod
- Tretman: clorhexidine kirurška tekućina - dezinficijens i nitrofurazone mast - tijekom 5 dana
- Ponovna kontrola za najmanje 10 dana

Spolne zaraze



Fig. 10-5 Uterus; cow. Venereal campylobacteriosis.
(Registry Acc. No. 7491, International Registry of Reproductive Pathology, Urbana, Illinois.)

Spolne zaraze




Fig. 10-7 Uterus; cow. Trichomoniasis.
(Courtesy K McEntee, Ithaca, New York.)



Fig. 10-12 Uterus; sow. Mixed flora endometritis.

CEH - pijometra kompleks

- CEH - cistična endometrijalna hiperplazija
- Endokrini poremećaj - opasan po život
- Kod kuja srednje i starije dobi
- Bez potomstva
- Hormonalno tretirane
- Razvije se tijekom lutealne faze
- Najčešće se ispoljava 30 - 45 dana nakon estrusa






Fig. 10-19 Uterus; bitch. Cystic endometrial hyperplasia-pyometra complex.



Fig. 10-21 Ovary; bitch. Retained corpora lutea.




Fig. 10-16 Uterus; bitch. Cystic endometrial hyperplasia-pyometra complex.




Fig. 10-17 Uterus; bitch. Cystic endometrial hyperplasia.





Fig. 10-18 Uterus; bitch. Cystic endometrial hyperplasia.



CEH - pijometra kompleks

Dijagnostika:

Anamneza:


- Tjerala se prije 30 - 60 dana
- Iscjedak iz stidnice - gnojni, sukrvavi, čokoladne boje
- Potištena, bolna u području abdomena, napet abdomen, teže se diže i liježe, povišena temperatura
- Slabiji apetit
- Ponekad izražena polidipsija

CEH - pijometra kompleks

Dijagnostika:

Objektivna

- RTG
- UZV
- Laboratorijske pretrage – leukocitoza



CEH - pijometra kompleks


Liječenje:


- Operativni zahvat - OVH
- Kod izuzetno vrijednih i mladih životinja terapija s PGF_{2α} i ATB












































Fig. 10-27 Uterus; cat. Pyometra.






Fig. 10-29 Uterus; mouse. Hydrometra.




Fig. 10-30 Uterus; mare. Mucometra.




Fig. 10-31 Uterus; mare. Endometrial polyps.




Fig. 10-39 Uterus; tiger (*Panthera tigris*). Adenocarcinoma. Large felidae.




Fig. 10-40 Uterus; tiger (*Panthera tigris*). Adenocarcinoma. Large felidae.



Fig. 10-44 Uterus; bitch. Adenocarcinoma.



Fig. 10-45 Uterus; cow. Leiomyoma.
Registry Acc. No. 4911, International Registry of Reproductive Pathology, Urbana, Illinois.)




Fig. 10-48 Uterus; bitch. Leiomyoma.




Fig. 10-49 Uterus; cow. Lymphosarcoma.



Fig. 10-47 Uterus; lion (*Panthera leo*). Leiomyoma.
Large felidae.



Fig. 10-50 Uterus; cow. Lymphosarcoma.



Fig. 10-52 Uterus; cow. Lymphosarcoma and pregnancy.




Fig. 10-53 Uterus; cow. Lymphosarcoma.
(Courtesy JM King, Ithaca, New York.)



Fig. 10-54 Uterus; pig. Lymphosarcoma.

References

1. Sheldon IM, Lewis GS, LeBlanc S, et al. Defining postpartum uterine disease in cattle. Theriogenology 2006 in press.
2. Smith BI, Donovan GA, Risco C, et al. Comparison of various antibiotic treatments for cows diagnosed with toxic puerperal metritis. J Dairy Sci 1998;81: 1555-62.
3. Overton MW, Sischo WM, Reynolds JP. Evaluation of effect of estradiol cypionate administration prophylactically to postparturient dairy cows at high risk for metritis. J Am Vet Med Assoc 2003; 223: 846-51.
4. Sheldon IM, Dobson H. Postpartum uterine health in cattle. Anim Reprod Sci 2004; 82/83: 295-306.
5. Lewis GS. Uterine health and disorders. J Dairy Sci 1997; 80: 984-94.
6. Sheldon IM, Rycroft AN, Zhou C. Association between postpartum pyrexia and uterine bacterial infection in dairy cattle. Vet Rec 2004; 154: 289-93.
7. Bondurant RH. Inflammation in the bovine female reproductive tract. J Dairy Sci 1999; 82: 101-10.
8. Dhaliwal GS, Murray RD, Woldehiwet Z. Some aspects of immunology of the bovine uterus related to treatments for endometritis. Anim Reprod Sci 2001; 67: 135-52.
9. Drillich M, Beetz O, Pfützner A, et al. Evaluation of a systemic antibiotic treatment of toxic puerperal metritis in dairy cows. J Dairy Sci 2001; 84: 2010-7.
10. Risco CA, Hernandez J. Comparison of ceftiofur hydrochloride and estradiol cypionate for metritis prevention and reproductive performance in dairy cows affected with retained fetal membranes. Theriogenology 2003; 60: 47-58.
11. Drillich M, Mahlstedt M, Reichert U, et al. Strategies to improve the therapy of retained fetal membranes in dairy cows. J Dairy Sci 2006 in press.
12. Okker H, Schmitt EJ, Vos PLAM, et al. Pharmacokinetics of ceftiofur in plasma and uterine secretions and tissues after suctaneous postpartum administration in lactating dairy cows. J Vet Pharmacol Ther 2002; 25: 33-8.
13. Königsson K, Gustafsson H, Gunnarsson A, et al. Clinical and bacteriological aspects on the use of Oxytetracycline and flunixin in primiparous cows with induced retained placenta and post-partal endometritis. Reprod Dom Anim 2001; 36: 247-56.
14. LeBlanc SJ, Duffield TF, Leslie KE, et al. Defining and diagnosing postpartum clinical endometritis and its impact on reproductive performance in dairy cows. J Dairy Sci 2002; 85: 2223-36.
15. Wittenbrink MM, Schoon HA, Schoon D, et al. Endometritis in cattle experimentally induced by Chlamydia psittaci. J Vet Med B 1993; 40: 437-50.
16. Bonnett BN, Martin SW, Meek AH. Associations of clinical findings, bacteriological and histological results of endometrial biopsy with reproductive performance of postpartum dairy cows. Prev Vet Med 193; 15: 205-20.
17. Williams EJ, Fischer DP, Pfeiffer DU, et al. Clinical evaluation of postpartum vaginal mucus reflects uterine bacterial infection and the immune response in cattle. Theriogenology 2005; 63: 101-17.
18. Falkenberg U, Heuwieser W. Influence of time of initiation of a prostaglandin F2alpha protocol in dairy cows with puerperal endometritis. Dtsch Tierarztl Wochenschr 2005; 112: 252-6.
19. Knutti B, Busato A, Küpfer U. Reproductive efficiency of cows with endometritis after treatment with intrauterine infusions or prostaglandin injections or no treatment. J Vet Med A 2000; 47: 609-15.
20. McDougall S. Effect of intrauterine antibiotic treatment on reproductive performance of dairy cows following periparturient disease. N Z Vet J 2001; 49: 150-8.
21. LeBlanc SJ, Duffield TF, Leslie KE, et al. The effect of treatment of clinical endometritis on reproductive performance in dairy cows. J Dairy Sci 2002; 85: 2237-49.
22. Heuwieser W, Tenhagen BA, Tischer M, et al. Effect of three programmes for the treatment of endometritis on the reproductive performance of a dairy herd. Vet Rec 2000; 146: 338-41.
23. Raab D, Drillich M, Heuwieser W. Diagnosis of subclinical endometritis and its effect on reproductive performance. In: 36th Annual conference of the American Association of Bovine Practitioners. Columbus, Ohio/USA, 2003: 166.
24. Kasimanickam R, Duffield TF, Foster RA, et al. Endometrial cytology and ultrasonography for the detection of subclinical endometritis in postpartum dairy cows. Theriogenology 2004; 62: 9-23.
25. Kasimanickam R, Duffield TF, Foster RA, et al. The effect of a single administration of cephalopirin or cloprostestol on the reproductive performance of dairy cows with subclinical endometritis. Theriogenology 2005; 63: 818-30.
26. Drillich M, Wittke M, Tenhagen BA, et al. Treatment of chronic endometritis in dairy cows with cephalopirin, tiaprost or a combination of both. Tierärztl Prax G 2005, 33: 404-10.
27. Nakao T, Moriyoshi M, Kawata K. Effect of postpartum intrauterine treatment with 2 % polyvinyl-pyrrolidone-iodine solution on reproductive efficiency in cows. Theriogenology 1988; 30: 1033-43.
28. Drillich M, Raab D, Wittke M, et al. Treatment of chronic endometritis in dairy cows with an intrauterine application of enzymes. A field trial. Theriogenology 2005; 63: 1811-23.
29. Lenz M, Drillich M, Heuwieser W. Ultrasound

Buergelt, CD (1997): *Color Atlas of Reproductive Pathology of Domestic Animals*, Mosby-Year Book Inc. 112-132.

Arthur, GH, DE Noakes and H. Pearson (2001): *Veterinary Reproduction & Obstetrics*, W.B. Saunders Comp.

Bearden HJ, and JW Fuquay (1992): *Applied Animal Reproduction*, Prentice Hall Inc. 262-263, 326.

Noakes, DE (1997): *Fertility and Obstetrics in Cattle*, Blackwell Science LTD. 106-109.

The Merck Veterinary Manual (2005): Merck & Co LTD, 1130-1134.

Youngquist, RS (1997): *Current Therapy in Large Animal Theriogenology*, W.B. Saunders Comp. 335 - 340.

Hafez, B and ESE Hafez (2000): *Reproduction in Farm Animals*, Lippincott Williams & Wilkins, 277 - 278.

Korisni linkovi

- <http://www.rvc.ac.uk>
- <http://www.vet.cornell.edu>
- <http://www.vetmed.wisc.edu/home>
- <http://www.vet.ohio-state.edu>
- <http://www.vetmed.fu-berlin.de>
- http://www.usask.ca/wcvm/links/intl_schools.php

Hvala, još koje pitanje?

